

Average Degree-Eccentricity Energy of Graphs

Veena Mathad, Shadi Ibrahim Khalaf, Sultan Senan Mahde
and Ivan Gutman*

Abstract

The concept of average degree-eccentricity matrix $ADE(G)$ of a connected graph G is introduced. Some coefficients of the characteristic polynomial of $ADE(G)$ are obtained, as well as a bound for the eigenvalues of $ADE(G)$. We also introduce the average degree-eccentricity graph energy and establish bounds for it.

Keywords: Average degree-eccentricity matrix, average degree-eccentricity eigenvalue, average degree-eccentricity energy.

2010 Mathematics Subject Classification: 05C50.

How to cite this article

V. Mathad, S. I. Khalaf, S. S. Mahde, I. Gutman, Average degree-eccentricity energy of graphs, *Math. Interdisc. Res.* **3** (2018) 45-54.

1. Introduction

Throughout this paper, all graphs are assumed to be simple, finite and connected. Let $G = (V, E)$ be such a graph, with vertex set \mathbf{V} and edge set \mathbf{E} . If $|\mathbf{V}| = p$ and $|\mathbf{E}| = q$, then G is said to be a (p, q) -graph. The degree of a vertex v , denoted by $d(v)$, is the number of edges of G incident with v . The distance $d(u, v)$ between two vertices u and v in a graph G is the length of a shortest path connecting them. For a vertex v of G , the eccentricity of v is $e(v) = \max\{d(v, u), u \in \mathbf{V}(G)\}$. For additional graph-theoretical terminologies we refer to [8].

The adjacency matrix of G , $\mathbf{A}(G) = (a_{ij})$ is a $p \times p$ matrix, such that $a_{ij} = 1$ if $v_i v_j \in \mathbf{E}$ and $a_{ij} = 0$ otherwise. The energy of G , denoted by $E(G)$, is defined as

$$E(G) = \sum_{i=1}^p |x_i| \tag{1}$$

*Corresponding author (E-mail: gutman@kg.ac.rs)

Academic Editor: Ali Reza Ashrafi

Received 13 February 2018, Accepted 20 February 2018

DOI: 10.22052/mir.2018.119231.1090



where x_1, x_2, \dots, x_p are the eigenvalues of $\mathbf{A}(G)$. This concept was introduced almost 40 years ago [5] and has been extensively investigated [2, 6, 7, 10]. Eventually, numerous other graph energies have been invented, based on eigenvalues of matrices different from the adjacency matrix; for more details see [1, 6, 7, 9, 11, 13–17] and the references cited therein.

One of these graph energies is the *sum-eccentricity energy* [15, 17], based on the eigenvalues of the *sum-eccentricity matrix* \mathbf{SE} , whose elements are equal defined as

$$se_{ij} = \begin{cases} e(v_i) + e(v_j) & \text{if } v_i v_j \in \mathbf{E} \\ 0 & \text{otherwise.} \end{cases} \quad (2)$$

Another recently introduced graph energy is the *first Zagreb energy* [9], based on the eigenvalues of the *first Zagreb matrix* \mathbf{ZG} , whose elements are defined as

$$zg_{ij} = \begin{cases} d(v_i) + d(v_j) & \text{if } v_i v_j \in \mathbf{E} \\ 0 & \text{otherwise.} \end{cases} \quad (3)$$

In this article, we introduce the concept of *average degree-eccentricity matrix* \mathbf{ADE} .

Definition 1.1. Let $G = (V, E)$ be a simple connected graph with p vertices v_1, v_2, \dots, v_p and let d_i and $e(v_i)$ be, respectively, the degree and eccentricity of v_i , $i = 1, 2, \dots, p$. Then the average degree-eccentricity matrix $\mathbf{ADE} = \mathbf{ADE}(G)$ of G is the $p \times p$ matrix whose elements are given by

$$m_{ij} = \begin{cases} \frac{1}{4}[d(v_i) + d(v_j) + e(v_i) + e(v_j)] & \text{if } v_i v_j \in \mathbf{E} \\ 0 & \text{otherwise.} \end{cases} \quad (4)$$

Bearing in mind Equations (2) and (3), we see that \mathbf{ADE} is conceived as a linear combination of the sum-eccentricity and Zagreb matrices, i.e.,

$$\mathbf{ADE} = \frac{1}{4}[\mathbf{SE} + \mathbf{ZG}].$$

The eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_p$ of $\mathbf{ADE}(G)$ form the average degree-eccentricity spectrum or the \mathbf{ADE} -spectrum of G . As usual, the \mathbf{ADE} -spectrum of G with n_i -fold degenerate eigenvalues λ_i is written as

$$S_p(G) = \{(\lambda_1)^{n_1}, (\lambda_2)^{n_2}, \dots, (\lambda_p)^{n_p}\}.$$

\mathbf{ADE} is a real symmetric matrix. Therefore, its eigenvalues are real numbers, and $\sum_{i=1}^p \lambda_i = 0$.

The following result will be useful in the proof of our results.

Theorem 1.2. [3] (*Gershgorin's Theorem*) Every eigenvalue λ of a $p \times p$ matrix $M = (m_{ij})$ satisfies:

$$|\lambda - m_{ii}| \leq \sum_{\substack{j=1 \\ j \neq i}}^p |m_{ij}|.$$

Corollary 1.3. [4] (*Hadamard's Inequality*) If the entries of a $p \times p$ matrix M are bounded by B , then $|\det(M)| \leq B^p p^{p/2}$.

2. Average Degree-Eccentricity Energy

Definition 2.1. The average degree-eccentricity energy $E_{ade}(G)$ of a graph G is

$$E_{ade}(G) = \sum_{i=1}^p |\lambda_i|. \quad (5)$$

Evidently, the average degree-eccentricity energy is defined in analogy to the ordinary graph energy, Equation (1).

Example 2.2. For a graph G_1 in Figure 1,




Figure 1: G_1 .

the average degree-eccentricity matrix of G_1 is

$$\mathbf{ADE}(G_1) = \begin{bmatrix} 0 & \frac{9}{4} & 0 & 0 & 0 \\ \frac{9}{4} & 0 & \frac{9}{4} & 0 & \frac{9}{4} \\ 0 & \frac{9}{4} & 0 & \frac{9}{4} & 0 \\ 0 & 0 & \frac{9}{4} & 0 & \frac{9}{4} \\ 0 & \frac{9}{4} & 0 & \frac{9}{4} & 0 \end{bmatrix}$$

The characteristic polynomial of $\mathbf{ADE}(G_1)$ is, $P(G_1, \lambda) = |\lambda I_p - \mathbf{ADE}(G_1)| = \lambda^5 - \frac{405}{16}\lambda^3 + \frac{6561}{128}\lambda$ and the average degree-eccentricity eigenvalues of G_1 are $\lambda_1 \approx 4.8, \lambda_2 \approx 1.5, \lambda_3 = 0, \lambda_4 \approx -1.5, \lambda_5 \approx -4.8$. Then the average degree-eccentricity energy of G_1 is $E_{ade}(G_1) = 4.8 + 1.5 + 1.5 + 4.8 = 12.6$.

We now calculate the coefficient c_i of λ^{p-i} ($i = 0, 1, 2, p$) in the characteristic polynomial of the average degree-eccentricity matrix $\mathbf{ADE}(G)$. Clearly $c_0 = 1$, $c_1 = \text{trace}(\mathbf{ADE}(G)) = 0$. Now

$$c_2 = \sum_{1 \leq i < j \leq p} \begin{vmatrix} 0 & m_{ij} \\ m_{ji} & 0 \end{vmatrix} = \sum_{1 \leq i < j \leq p} -m_{ij}^2.$$

In view of Equation (4) we get

$$c_2 = - \sum_{v_i v_j \in \mathbf{E}} \left[\frac{d(v_i) + e(v_i) + d(v_j) + e(v_j)}{4} \right]^2.$$

For c_3 we have

$$c_3 = (-1)^3 \sum_{1 \leq i < j < r \leq n} \begin{vmatrix} m_{ii} & m_{ij} & m_{ir} \\ m_{ji} & m_{jj} & m_{jr} \\ m_{ri} & m_{rj} & m_{rr} \end{vmatrix}.$$

The number of non-zero terms in the above sum is equal to the number of triangles in G . Therefore, $c_3 = 0$ if G has no triangle.

Finally, $c_p = \det(\mathbf{ADE}(G))$.

Lemma 2.3. *Let G be a connected (p, q) -graph and $uv \in \mathbf{E}$. Then*

$$\frac{1}{4}[d(u) + d(v) + e(u) + e(v)] \leq \frac{p}{2}. \quad (6)$$

Equality in (6) holds for all $uv \in \mathbf{E}$ only if $G \cong K_p$.

Proof. Without loss of generality, we may assume that $e(u) \leq e(v)$. So, we have

$$\begin{aligned} d(u) + d(v) + e(u) + e(v) &\leq d(u) + d(v) + 2e(v) \\ &\leq d(u) + d(v) + 2[p - (d(u) + d(v)) + 1] \\ &= 2p - (d(u) + d(v)) + 2 \leq 2p. \end{aligned}$$

If $G \cong K_p$, then for any $uv \in \mathbf{E}$ we have $d(u) = d(v) = p - 1$ and $e(u) = e(v) = 1$, implying that the left-hand side of (6) is equal to $p/2$. For all other (connected) graphs, for some $uv \in \mathbf{E}$ the inequality in (6) will be strict. \square

Lemma 2.4. *Let G be a connected (p, q) -graph. Then*

$$\text{trace } \mathbf{ADE}^2(G) \leq \text{trace } \mathbf{ADE}^2(K_p) = \frac{(p-1)p^3}{4}. \quad (7)$$

Equality in (7) holds if and only if $G \cong K_p$.

Proof. Since

$$\mathbf{ADE}(K_p)_{ij} = \begin{cases} \frac{p}{2} & \text{if } v_i v_j \in \mathbf{E} \\ 0 & \text{otherwise} \end{cases}$$

we get that for $i \neq j$,

$$\mathbf{ADE}^2(K_p)_{ij} = (p-2) \left(\frac{p}{2}\right)^2$$

whereas for $i = j$,

$$\mathbf{ADE}^2(K_p)_{ii} = (p-1) \left(\frac{p}{2}\right)^2$$

implying that

$$\text{trace } \mathbf{ADE}^2(K_p) = p \times (p-1) \left(\frac{p}{2}\right)^2 = \frac{(p-1)p^3}{4}.$$

Bearing in mind Lemma 2.3 and formula (4), we immediately see that $\mathbf{ADE}(G)_{ij} \leq \mathbf{ADE}(K_p)_{ij}$, and that if $G \not\cong K_p$, then the inequality is strict for at least some of ij . Consequently, inequality (7) holds. \square

Theorem 2.5. *For any (p, q) -graph, with average degree-eccentricity eigenvalue λ_j ,*

$$|\lambda_j| \leq \frac{p(p-1)}{2}. \quad (8)$$

Proof. By Lemma 2.4, the trace of $\mathbf{ADE}^2(K_p)$ is equal to $\frac{(p-1)p^3}{4}$. Then for any (p, q) -graph G with average degree-eccentricity eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_p$, we have $\sum_{i=1}^p |\lambda_i|^2 \leq \frac{(p-1)p^3}{4}$. By the Cauchy–Schwarz inequality,

$$\left(\sum_{\substack{i=1 \\ i \neq j}}^p \lambda_i \right)^2 = (p-1) \sum_{\substack{i=1 \\ i \neq j}}^p \lambda_i^2.$$

Since $\sum_{i=1}^p \lambda_i^2 = -2c_2$ and $\sum_{i=1}^p \lambda_i = 0$, we get

$$\lambda_j^2 \leq (p-1) \left[\frac{(p-1)p^3}{4} - \lambda_j^2 \right]$$

which implies (8). \square

Proposition 2.6. *Let G be a graph of order p , and average degree-eccentricity eigenvalue λ_i . Then*

$$\prod_{i=1}^p |\lambda_i| \leq \left(\frac{p}{2}\right)^p p^{p/2}.$$

Proof. By Corollary 1.3 and by the definition of **ADE**, setting $B = p/2$. \square

Theorem 2.7. *Let G be a (p, q) -graph. Then*

$$E_{ade}(G) \leq \frac{(p-1)p^2}{2}.$$

Proof. By Gershgorin's Theorem and Lemma 2.3, we have

$$\begin{aligned} E_{ade}(G) &= \sum_{i=1}^p |\lambda_i| = \sum_{i=1}^p |\lambda_i - 0| \leq \sum_{i=1}^p \sum_{\substack{j=1 \\ j \neq i}}^p m_{ij} \\ &\leq \sum_{i=1}^p \sum_{\substack{j=1 \\ j \neq i}}^p \frac{p}{2} = \frac{(p-1)p^2}{2}. \end{aligned}$$

\square

Theorem 2.8. *Let G be a connected (p, q) -graph. Then*

$$E_{ade}(G) \geq \sqrt{2(q|\det(\mathbf{ADE})|^{2/p} - c_2)}.$$

Proof.

$$E_{ade}(G)^2 = \left(\sum_{i=1}^p |\lambda_i| \right)^2 = \sum_{i=1}^p \lambda_i^2 + \sum_{\substack{i=1 \\ i \neq j}}^p |\lambda_i| |\lambda_j| = -2c_2 + \sum_{\substack{i=1 \\ i \neq j}}^p |\lambda_i| |\lambda_j|.$$

From relation between the arithmetic and geometric means, we get

$$\begin{aligned} \sum_{\substack{i=1 \\ i \neq j}}^p |\lambda_i| |\lambda_j| &\geq p(p-1) \left(\prod_{\substack{i=1 \\ i \neq j}}^p |\lambda_i| |\lambda_j| \right)^{\frac{1}{p(p-1)}} = p(p-1) \left(\prod_{\substack{i=1 \\ i \neq j}}^p |\lambda_i|^{2(p-1)} \right)^{\frac{1}{p(p-1)}} \\ &= p(p-1) \left(\prod_{\substack{i=1 \\ i \neq j}}^p |\lambda_i| \right)^{2/p} \geq 2q \left(\prod_{\substack{i=1 \\ i \neq j}}^p |\lambda_i| \right)^{2/p} = 2q |\det(\mathbf{ADE})|^{2/p}. \end{aligned}$$

Then

$$E_{ade}(G)^2 \geq 2q |\det(\mathbf{ADE})|^{2/p} - 2c_2 = 2[q |\det(\mathbf{ADE})|^{2/p} - c_2]$$

and finally,

$$E_{ade}(G) \geq \sqrt{2[q |\det(\mathbf{ADE})|^{2/p} - c_2]}.$$

\square

Note that Theorem 2.8 and its proof are just a replica of the classical McClelland inequality for ordinary graph energy [12].

Corollary 2.9. *Let G be a connected (p, q) -graph. Then*

$$\sqrt{2(q|\det(\mathbf{ADE})|^{2/p} - c_2)} \leq E_{ade}(G) \leq \frac{(p-1)p^2}{2}.$$

3. Average Degree-Energy of Some Classes of Graphs

In this section, we compute the average degree-eccentricity energies of some well-known graphs.

Example 3.1. Let G be a complete graph K_p . Then $S_p(\mathbf{ADE}(K_p)) = \{(\frac{p}{2})^{p-1}, ((p-1)(\frac{p}{2}))^1\}$ and $E_{ade}(K_p) = 2(p-1)(\frac{p}{2})$.

Proof. Let G be the complete graph K_p . Then

$$\begin{aligned} |\lambda I - \mathbf{ADE}(K_p)| &= \begin{vmatrix} \lambda & -\frac{p}{2} & -\frac{p}{2} & \cdots & -\frac{p}{2} \\ -\frac{p}{2} & \lambda & -\frac{p}{2} & \cdots & -\frac{p}{2} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ -\frac{p}{2} & -\frac{p}{2} & -\frac{p}{2} & \cdots & \lambda \end{vmatrix} \\ &= \left(\lambda + \frac{p}{2}\right)^{p-1} \begin{vmatrix} \lambda & -\frac{p}{2} & -\frac{p}{2} & \cdots & -\frac{p}{2} \\ -1 & 1 & 0 & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ -1 & 0 & 0 & \cdots & 1 \end{vmatrix} \\ &= \left(\lambda + \frac{p}{2}\right)^{p-1} \left[\lambda - (p-1)\frac{p}{2}\right]. \end{aligned}$$

Then the average degree-eccentricity energy of the complete graph is

$$E_{ade}(K_p) = 2(p-1)\frac{p}{2}.$$

□

Example 3.2. Let G be a complete bipartite graph $K_{m,n}$, $m, n \geq 2$. Then

$$S_p(\mathbf{ADE}(K_{m,n})) = \left\{ \left(\frac{p+4}{4}\sqrt{mn}\right)^1, (0)^{p-2}, \left(-\frac{p+4}{4}\sqrt{mn}\right)^1 \right\} \quad (9)$$

and

$$E_{ade}(K_{m,n}) = \frac{p+4}{2}\sqrt{mn}. \quad (10)$$

Proof. Let the vertex set of $K_{m,n}$ be $V = \{v_1, v_2, \dots, v_m, u_1, u_2, \dots, u_n\}$. Then, $p = m + n$, $q = mn$, and

$$\begin{aligned}
 |\lambda I - \mathbf{ADE}(K_{m,n})| &= \begin{vmatrix} \lambda & 0 & \cdots & 0 & -\frac{p+4}{4} & \cdots & -\frac{p+4}{4} \\ 0 & \lambda & \cdots & 0 & -\frac{p+4}{4} & \cdots & -\frac{p+4}{4} \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & \cdots & 0 & \lambda & -\frac{p+4}{4} & \cdots & -\frac{p+4}{4} \\ -\frac{p+4}{4} & \cdots & -\frac{p+4}{4} & 0 & \lambda & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ -\frac{p+4}{4} & \cdots & -\frac{p+4}{4} & 0 & \cdots & 0 & \lambda \end{vmatrix} \\
 &= \lambda^p - \left(\frac{p+4}{4}\right)^2 (mn)\lambda^{p-2} = \lambda^{p-2} \left[\lambda^2 - \left(\frac{p+4}{4}\right)^2 (mn) \right].
 \end{aligned}$$

Then, $\lambda^{p-2} \left[\lambda^2 - \left(\frac{p+4}{4}\right)^2 (mn) \right] = 0$ implies $\lambda^{p-2} = 0$, or $\lambda^2 = \left(\frac{p+4}{4}\right)^2 (mn)$, resulting in (9) and (10). \square

Example 3.3. For the star graph $K_{1,p-1}$,

$$S_p(\mathbf{ADE}(K_{1,p-1})) = \left\{ \left(\frac{p+3}{4}\sqrt{p-1}\right)^1, (0)^{p-2}, \left(-\frac{p+3}{4}\sqrt{p-1}\right)^1 \right\}$$

and

$$E_{ade}(K_{1,p-1}) = \frac{p+3}{2}\sqrt{p-1}.$$

Acknowledgement. The authors thank the anonymous referee for useful comments and suggestion.

Conflicts of Interest. The authors declare that they have no conflicts of interest.

References

- [1] C. Adiga, M. Smitha, On maximum degree energy of a graph, *Int. J. Contemp. Math. Sci.* **4** (2009) 385–396.
- [2] R. Balakrishnan, The energy of a graph, *Linear Algebra Appl.* **387** (2004) 287–295.
- [3] H. E. Bell, Gerschgorin's theorem and the zeros of polynomials, *Am. Math. Monthly* **72** (1965) 292–295.
- [4] D. J. H. Garling, *Inequalities – A Journey Into Linear Analysis*, Cambridge Univ. Press, Cambridge, 2007.

-
- [5] I. Gutman, The energy of a graph, *Ber. Math.-Statist. Sect. Forsch. Graz* **103** (1978) 1–22.
- [6] I. Gutman, B. Furtula, Survey of graph energies, *Math. Interdisc. Res.* **2** (2017) 85–129.
- [7] I. Gutman, B. Furtula, The total π -electron energy saga, *Croat. Chem. Acta* **90** (2017) 359–368.
- [8] F. Harary, *Graph Theory*, Addison Wesley, Reading, 1969.
- [9] N. Jafari Rad, A. Jahanbani, I. Gutman, Zagreb energy and Zagreb Estrada index of graphs, *MATCH Commun. Math. Comput. Chem.* **79** (2018) 371–386.
- [10] X. Li, Y. Shi, I. Gutman, *Graph Energy*, Springer, New York, 2012.
- [11] V. Mathad, S. S. Mahde, The minimum hub energy of a graph, *Palest. J. Math.* **6** (2017) 247–256.
- [12] B. J. McClelland, Properties of the latent roots of a matrix: The estimation of π -electron energies, *J. Chem. Phys.* **54** (1971) 640–643.
- [13] M. A. Naji, N. D. Soner, The maximum eccentricity energy of a graph, *Int. J. Sci. Engin. Res.* **7** (2016) 5–13.
- [14] H. S. Ramane, I. Gutman, J. B. Patil, R. B. Jummannaver, Seidel signless Laplacian energy of graphs, *Math. Interdisc. Res.* **2** (2017) 181–192.
- [15] D. S. Revankar, M. M. Patil, H. S. Ramane, On eccentricity sum eigenvalue and eccentricity sum energy of a graph, *Ann. Pure Appl. Math.* **13** (2017) 125–130.
- [16] B. Sharada, M. I. Sowaity, I. Gutman, Laplacian sum-eccentricity energy of a graph, *Math. Interdisc. Res.* **2** (2017) 209–219.
- [17] M. I. Sowaity, B. Sharada, The sum-eccentricity energy of a graph, *Int. J. Rec. Innovat. Trends Comput. Commun.* **5** (2017) 293–304.

Veena Mathad
Department of Studies in Mathematics,
University of Mysore, Manasagangotri,
Mysuru – 570 006, India
E-mail: veena_mathad@rediffmail.com

Shadi Ibrahim Khalaf
Department of Studies in Mathematics,

University of Mysore, Manasagangotri,
Mysuru – 570 006, India
E-mail: shadikhalaf1989@hotmail.com

Sultan Senan Mahde
Department of Studies in Mathematics,
University of Mysore, Manasagangotri,
Mysuru – 570 006, India
E-mail: sultan.mahde@gmail.com

Ivan Gutman
Faculty of Science,
University of Kragujevac,
P. O. Box 60, 34000 Kragujevac, Serbia
E-mail: gutman@kg.ac.rs