

## Seidel Signless Laplacian Energy of Graphs

*Harishchandra S. Ramane\**, *Ivan Gutman*,  
*Jayashri B. Patil and Raju B. Jummanner*

### Abstract

Let  $S(G)$  be the Seidel matrix of a graph  $G$  of order  $n$  and let  $D_S(G) = \text{diag}(n-1-2d_1, n-1-2d_2, \dots, n-1-2d_n)$  be the diagonal matrix with  $d_i$  denoting the degree of a vertex  $v_i$  in  $G$ . The Seidel Laplacian matrix of  $G$  is defined as  $SL(G) = D_S(G) - S(G)$  and the Seidel signless Laplacian matrix as  $SL^+(G) = D_S(G) + S(G)$ . The Seidel signless Laplacian energy  $E_{SL^+}(G)$  is defined as the sum of the absolute deviations of the eigenvalues of  $SL^+(G)$  from their mean. In this paper, we establish the main properties of the eigenvalues of  $SL^+(G)$  and of  $E_{SL^+}(G)$ .

**Keywords:** Seidel Laplacian eigenvalues, Seidel Laplacian energy, Seidel signless Laplacian matrix, Seidel signless Laplacian eigenvalues, Seidel signless Laplacian energy.

2010 Mathematics Subject Classification: 05C50.

## 1. Introduction

Let  $G$  be a simple, undirected graph with  $n$  vertices and  $m$  edges. We say that  $G$  is an  $(n, m)$ -graph.

Let  $v_1, v_2, \dots, v_n$  be the vertices of  $G$ . The degree of a vertex  $v_i$  is the number of edges incident to it and is denoted by  $d_i$ . If  $d_i = r$  for all  $i = 1, 2, \dots, n$ , then  $G$  is said to be an  $r$ -regular graph.

By  $\overline{G}$  will be denoted the complement of the graph  $G$ .

The *adjacency matrix* of a graph  $G$  is the square matrix  $A(G) = (a_{ij})$ , in which  $a_{ij} = 1$  if  $v_i$  is adjacent to  $v_j$  and  $a_{ij} = 0$  otherwise. If  $\lambda_1, \lambda_2, \dots, \lambda_n$  are

---

\*Corresponding author (E-mail: [hsramane@yahoo.com](mailto:hsramane@yahoo.com))  
Academic Editor: Ali Reza Ashrafi  
Received 20 October 2017, Accepted 6 November 2017  
DOI: 10.22052/mir.2017.101641.1081

the eigenvalues of  $A(G)$ , then the *ordinary energy* of a graph  $G$  is defined as [14,19]

$$E_A(G) = \sum_{i=1}^n |\lambda_i|. \quad (1)$$

The spectral graph theory based on the eigenvalues of the adjacency matrix is the most extensively elaborated part of algebraic graph theory [4, 6]. Also the graph energy, based on the eigenvalues of the adjacency matrix, attracted much attention and has been studied to a great extent [19]. Recently, a number of other graph energies have been introduced; for details see the recent survey [16]. Among these, the Laplacian and Seidel energies are of importance for the following considerations.

Let  $D(G) = \text{diag}(d_1, d_2, \dots, d_n)$  be the diagonal degree matrix. The *Laplacian matrix* of  $G$  is defined as  $L(G) = D(G) - A(G)$ . This matrix and its spectral properties have been extensively studied in the past [12, 13, 20–22]. Let  $\mu_1, \mu_2, \dots, \mu_n$  be the eigenvalues of  $L(G)$ . Then the *Laplacian energy* of  $G$  is defined as [17]

$$E_L(G) = \sum_{i=1}^n \left| \mu_i - \frac{2m}{n} \right|. \quad (2)$$

In 2007, Cvetković et al. [5] noticed that if the definition of the Laplacian matrix is changed into  $D(G) + A(G)$ , then interesting and non-trivial spectral properties are obtained. Eventually, an entire theory of the so-called signless Laplacian spectra has been developed [7–9].

Thus, the *signless Laplacian matrix* of  $G$  is defined as  $L^+(G) = D(G) + A(G)$ . Let  $\mu_1^+, \mu_2^+, \dots, \mu_n^+$  be its eigenvalues. Then the *signless Laplacian energy* of  $G$  is defined as [1]

$$E_{L^+}(G) = \sum_{i=1}^n \left| \mu_i^+ - \frac{2m}{n} \right|. \quad (3)$$

The *Seidel matrix* of a graph  $G$  is the  $n \times n$  real symmetric matrix  $S(G) = (s_{ij})$ , where  $s_{ij} = -1$  if the vertices  $v_i$  and  $v_j$  are adjacent,  $s_{ij} = 1$  if the vertices  $v_i$  and  $v_j$  are not adjacent, and  $s_{ij} = 0$  if  $i = j$ . It is easy to see that  $S(G) = A(\bar{G}) - A(G)$ , where  $\bar{G}$  is the complement of  $G$ .

The eigenvalues of the Seidel matrix, labeled as  $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_n$ , are said to be the *Seidel eigenvalues* of  $G$  and form the Seidel spectrum. In analogy to Eq. (1), the *Seidel energy* of a graph  $G$  is defined as [18]

$$E_S(G) = \sum_{i=1}^n |\sigma_i|. \quad (4)$$

More results on Seidel energy are reported in [11, 23, 24, 27–29].

The Seidel Laplacian matrix of a graph was introduced in [30], where the main properties of its eigenvalues and of Seidel Laplacian energy were established.

Let  $D_S(G) = \text{diag}(n-1-2d_1, n-1-2d_2, \dots, n-1-2d_n)$  be a diagonal matrix in which  $d_i$  stands for degree of a vertex  $v_i$  in  $G$ . Then, in analogy with the ordinary Laplacian matrix, the *Seidel Laplacian matrix* of  $G$  is defined as

$$SL(G) = D_S(G) - S(G).$$

Note that  $D_S(G) = D(\overline{G}) - D(G)$  and  $SL(G) = L(\overline{G}) - L(G)$ .

Let  $\sigma_1^L, \sigma_2^L, \dots, \sigma_n^L$ , be the eigenvalues of  $SL(G)$ . In analogy to Eq. (2), the *Seidel Laplacian energy* of  $G$  is defined as [30]

$$E_{SL}(G) = \sum_{i=1}^n \left| \sigma_i^L - \frac{n(n-1) - 4m}{n} \right|.$$

Bearing in mind the concept of signless Laplacian matrix, we introduce here the Seidel signless Laplacian matrix of a graph and study the basic properties of its eigenvalues and energy.

Thus, the *Seidel signless Laplacian matrix* of  $G$  is defined as

$$SL^+(G) = D_S(G) + S(G).$$

In the case of Laplacian matrices, the adjective “signless” is fully appropriate, since no element of  $L^+(G)$  is negative-valued. In the case of Seidel Laplacians, we use the same adjective by analogy, aware of the fact that all Seidel signless Laplacian matrices, with the exception of  $SL^+(K_n)$ , possess negative-valued elements, where  $K_n$  denotes the complete graph on  $n$  vertices.

Note that

$$\begin{aligned} SL^+(G) &= D(\overline{G}) - D(G) + A(\overline{G}) - A(G) \\ &= D(\overline{G}) + A(\overline{G}) - D(G) - A(G) \\ &= L^+(\overline{G}) - L^+(G). \end{aligned}$$

Let  $\sigma_1^{L^+}, \sigma_2^{L^+}, \dots, \sigma_n^{L^+}$ , be the eigenvalues of  $SL^+(G)$ . In analogy to Eq. (3), the *Seidel signless Laplacian energy* of  $G$  is defined as

$$E_{SL^+}(G) = \sum_{i=1}^n \left| \sigma_i^{L^+} - \frac{n(n-1) - 4m}{n} \right|.$$

If we introduce the auxiliary quantities

$$\xi_i = \sigma_i^{L^+} - \frac{n(n-1) - 4m}{n}, \quad i = 1, 2, \dots, n$$

then the expression for Seidel signless Laplacian energy becomes analogous to the formula for ordinary graph energy, Eq. (1), namely:

$$E_{SL^+}(G) = \sum_{i=1}^n |\xi_i|.$$

## 2. Seidel Signless Laplacian Eigenvalues

The adjacency eigenvalues, Laplacian eigenvalues and Seidel eigenvalues satisfy the relations

$$\begin{aligned} \sum_{i=1}^n \lambda_i &= 0 & \text{and} & & \sum_{i=1}^n \lambda_i^2 &= 2m; \\ \sum_{i=1}^n \mu_i &= 2m & \text{and} & & \sum_{i=1}^n \mu_i^2 &= 2m + Z_1(G); \\ \sum_{i=1}^n \sigma_i &= 0 & \text{and} & & \sum_{i=1}^n \sigma_i^2 &= n(n-1) \end{aligned}$$

where  $Z_1(G) = \sum_{i=1}^n d_i^2$  is the well-known graph invariant called first Zagreb index [3, 15].

The respective formulas for the Seidel Laplacian eigenvalues read [30]:

$$\sum_{i=1}^n \sigma_i^L = n(n-1) - 4m \quad \text{and} \quad \sum_{i=1}^n (\sigma_i^L)^2 = (n-1)(n^2 - 8m) + 4Z_1(G). \quad (5)$$

In the next lemma, we show that the Seidel signless Laplacian eigenvalues have properties fully analogous to those of the ordinary Seidel Laplacians, Eqs. (5).

**Lemma 2.1.** *Let  $G$  be an  $(n, m)$ -graph. Then the eigenvalues of the Seidel signless Laplacian matrix satisfy the relations:*

$$\sum_{i=1}^n \sigma_i^{L^+} = n(n-1) - 4m \quad \text{and} \quad \sum_{i=1}^n (\sigma_i^{L^+})^2 = (n-1)(n^2 - 8m) + 4Z_1(G).$$

*Proof.*

$$\sum_{i=1}^n \sigma_i^{L^+} = \text{trace}[SL^+(G)] = \sum_{i=1}^n (n-1 - 2d_i) = n(n-1) - 4m.$$

$$\begin{aligned} \sum_{i=1}^n (\sigma_i^{L^+})^2 &= \text{trace}[(SL^+(G))^2] \\ &= \sum_{i=1}^n [(n-1 - 2d_i)^2 + (n-1)] \\ &= n^2(n-1) - 8m(n-1) + 4Z_1(G) \\ &= (n-1)(n^2 - 8m) + 4Z_1(G). \end{aligned}$$

□

**Lemma 2.2.** Let  $\sigma_1^{L^+}, \sigma_2^{L^+}, \dots, \sigma_n^{L^+}$  be the eigenvalues of  $SL^+(G)$ . Let  $\xi_i = \sigma_i^{L^+} - \frac{n(n-1)-4m}{n}$ ,  $i = 1, 2, \dots, n$ . Then

$$\sum_{i=1}^n \xi_i = 0 \quad \text{and} \quad \sum_{i=1}^n \xi_i^2 = M$$

where  $M = n(n-1) + 4Z_1(G) - \frac{16m^2}{n}$ .

*Proof.*

$$\begin{aligned} \sum_{i=1}^n \xi_i &= \sum_{i=1}^n \left( \sigma_i^{L^+} - \frac{n(n-1)-4m}{n} \right) \\ &= \sum_{i=1}^n \sigma_i^{L^+} - [n(n-1) - 4m] = n(n-1) - 4m - [n(n-1) - 4m] = 0. \end{aligned}$$

$$\begin{aligned} \sum_{i=1}^n \xi_i^2 &= \sum_{i=1}^n \left( \sigma_i^{L^+} - \frac{n(n-1)-4m}{n} \right)^2 \\ &= \sum_{i=1}^n (\sigma_i^{L^+})^2 - 2 \left( \frac{n(n-1)-4m}{n} \right) \sum_{i=1}^n \sigma_i^{L^+} + \sum_{i=1}^n \left( \frac{n(n-1)-4m}{n} \right)^2 \\ &= (n-1)(n^2 - 8m) + 4Z_1(G) - 2 \left( \frac{n(n-1)-4m}{n} \right) (n(n-1) - 4m) \\ &\quad + n \left[ \frac{(n(n-1)-4m)^2}{n^2} \right] = n(n-1) + 4Z_1(G) - \frac{16m^2}{n}. \end{aligned}$$

□

As a consequence of Lemmas 2.1 and 2.2, the lower and upper bounds for the Seidel signless Laplacian energy stated below in Theorems 3.1, 3.2, 3.7–3.11 are fully analogous to the results earlier obtained for the ordinary Seidel Laplacian energy [30].

The next proposition follows from  $SL^+(G) = L^+(\overline{G}) - L^+(G) = -SL^+(\overline{G})$ .

**Proposition 2.3.** If  $\sigma_i^{L^+}$ ,  $i = 1, 2, \dots, n$ , are the Seidel signless Laplacian eigenvalues of  $G$ , then  $-\sigma_i^{L^+}$ ,  $i = 1, 2, \dots, n$ , are the Seidel signless Laplacian eigenvalues of  $\overline{G}$ .

**Theorem 2.4.** If  $\sigma_1, \sigma_2, \dots, \sigma_n$  are the Seidel eigenvalues of an  $r$ -regular graph  $G$ , then the Seidel signless Laplacian eigenvalues of  $G$  are  $n-1-2r+\sigma_i$ ,  $i = 1, 2, \dots, n$ .

*Proof.* Let the characteristic polynomial of the Seidel matrix  $S(G)$  be denoted by  $\phi_S(G, \lambda)$  and the characteristic polynomial of the Seidel signless Laplacian matrix  $SL^+(G)$  by  $\phi_{SL^+}(G, \lambda)$ .

If  $G$  is an  $r$ -regular graph, then  $SL^+(G) = (n - 1 - 2r)I + S(G)$ , where  $I$  is the identity matrix of order  $n$ . Therefore,

$$\begin{aligned}\phi_{SL^+}(G, \lambda) &= \det(\lambda I - SL^+(G)) \\ &= \det[(\lambda - n + 1 + 2r)I - S(G)] \\ &= \phi_S(G, \lambda - n + 1 + 2r).\end{aligned}$$

□

**Lemma 2.5.** *Let the notation be same as in Lemma 2.2. Then*

$$\left| \sum_{i < j} \xi_i \xi_j \right| = \frac{1}{2}M.$$

*Proof.* Since  $\sum_{i=1}^n \xi_i = 0$ , we get  $\sum_{i=1}^n \xi_i^2 = -2 \sum_{i < j} \xi_i \xi_j$ . Therefore

$$2 \left| \sum_{i < j} \xi_i \xi_j \right| = \sum_{i=1}^n \xi_i^2 = M.$$

□

### 3. Seidel Signless Laplacian Energy

**Theorem 3.1.** *Let  $G$  be an  $(n, m)$ -graph. Then*

$$E_{SL^+}(G) \leq \sqrt{n \left[ n(n-1) + 4Z_1(G) - \frac{16m^2}{n} \right]}. \quad (6)$$

*Proof.* The Cauchy–Schwartz inequality states that,

$$\left( \sum_{i=1}^n a_i b_i \right)^2 \leq \left( \sum_{i=1}^n a_i^2 \right) \left( \sum_{i=1}^n b_i^2 \right).$$

Set  $a_i = 1$  and  $b_i = |\xi_i|$ ,  $i = 1, 2, \dots, n$ . Then

$$\begin{aligned}\left( \sum_{i=1}^n |\xi_i| \right)^2 &\leq n \sum_{i=1}^n |\xi_i|^2 \\ [E_{SL^+}(G)]^2 &\leq n \left[ n(n-1) + 4Z_1(G) - \frac{16m^2}{n} \right]\end{aligned}$$

and inequality (6) follows. □

**Theorem 3.2.** *Let  $G$  be an  $(n, m)$ -graph. Then*

$$E_{SL^+}(G) \geq \sqrt{2 \left[ n(n-1) + 4Z_1(G) - \frac{16m^2}{n} \right]}.$$

*Proof.*  $E_{SL^+}(G) = \sum_{i=1}^n |\xi_i|$ . Therefore

$$\begin{aligned} [E_{SL^+}(G)]^2 &= \sum_{i=1}^n |\xi_i|^2 + 2 \sum_{1 \leq i < j \leq n} |\xi_i| |\xi_j| \geq \sum_{i=1}^n |\xi_i|^2 + 2 \left| \sum_{1 \leq i < j \leq n} \xi_i \xi_j \right| \\ &= M + M = 2 \left[ n(n-1) + 4Z_1(G) - \frac{16m^2}{n} \right]. \end{aligned}$$

□

**Theorem 3.3.** [25] *Suppose that  $a_i$  and  $b_i$ ,  $1 \leq i \leq n$ , are positive real numbers. Then*

$$\sum_{i=1}^n a_i^2 \sum_{i=1}^n b_i^2 \leq \frac{1}{4} \left( \sqrt{\frac{M_1 M_2}{m_1 m_2}} + \sqrt{\frac{m_1 m_2}{M_1 M_2}} \right)^2 \left( \sum_{i=1}^n a_i b_i \right)^2$$

where  $M_1 = \max_{1 \leq i \leq n} (a_i)$ ,  $M_2 = \max_{1 \leq i \leq n} (b_i)$ ,  $m_1 = \min_{1 \leq i \leq n} (a_i)$ , and  $m_2 = \min_{1 \leq i \leq n} (b_i)$ .

**Theorem 3.4.** [26] *Let  $a_i$  and  $b_i$ ,  $1 \leq i \leq n$ , be non-negative real numbers. Then*

$$\sum_{i=1}^n a_i^2 \sum_{i=1}^n b_i^2 - \left( \sum_{i=1}^n a_i b_i \right)^2 \leq \frac{n^2}{4} (M_1 M_2 - m_1 m_2)^2$$

where  $M_i$  and  $m_i$ ,  $i = 1, 2$ , are defined similarly as in Theorem 3.3.

**Theorem 3.5.** [10] *Suppose that  $a_i$  and  $b_i$ ,  $1 \leq i \leq n$ , are positive real numbers. Then*

$$\left| n \sum_{i=1}^n a_i b_i - \sum_{i=1}^n a_i \sum_{i=1}^n b_i \right| \leq \alpha(n)(A - a)(B - b)$$

where  $a, b, A$ , and  $B$  are real constants, such that for each  $i$ ,  $1 \leq i \leq n$ ,  $a \leq a_i \leq A$ , and  $b \leq b_i \leq B$ . In addition,  $\alpha(n) = n \lfloor \frac{n}{2} \rfloor \left( 1 - \frac{1}{n} \lfloor \frac{n}{2} \rfloor \right)$ .

**Theorem 3.6.** [2] *Let  $a_i$  and  $b_i$ ,  $1 \leq i \leq n$ , be non-negative real numbers. Then*

$$\sum_{i=1}^n b_i^2 + rR \sum_{i=1}^n a_i^2 \leq (r + R) \left( \sum_{i=1}^n a_i b_i \right)$$

where  $r$  and  $R$  are real constants, such that for each  $i$ ,  $1 \leq i \leq n$ ,  $r a_i \leq b_i \leq R a_i$  holds.

**Theorem 3.7.** Let  $G$  be an  $(n, m)$ -graph and let  $\xi_{max} = \max_{1 \leq i \leq n} |\xi_i|$  and  $\xi_{min} = \min_{1 \leq i \leq n} |\xi_i|$ . Then

$$E_{SL^+}(G) \geq \sqrt{nM - \frac{n^2}{4}(\xi_{max} - \xi_{min})^2}. \quad (7)$$

*Proof.* Applying Theorem 3.4 for  $a_i = 1$  and  $b_i = |\xi_i|$  we get

$$\begin{aligned} \sum_{i=1}^n 1^2 \sum_{i=1}^n |\xi_i|^2 - \left( \sum_{i=1}^n |\xi_i| \right)^2 &\leq \frac{n^2}{4}(\xi_{max} - \xi_{min})^2 \\ nM - (E_{SL^+}(G))^2 &\leq \frac{n^2}{4}(\xi_{max} - \xi_{min})^2 \end{aligned}$$

from which inequality (7) is straightforward.  $\square$

**Theorem 3.8.** Let  $G$ ,  $\xi_{max}$ , and  $\xi_{min}$  be same as in Theorem 3.7. Then

$$E_{SL^+}(G) \geq \frac{2\sqrt{nM \xi_{max} \xi_{min}}}{\xi_{max} + \xi_{min}}. \quad (8)$$

*Proof.* Applying Theorem 3.3 for  $a_i = 1$  and  $b_i = |\xi_i|$  we get

$$\begin{aligned} \sum_{i=1}^n 1^2 \sum_{i=1}^n |\xi_i|^2 &\leq \frac{1}{4} \left( \sqrt{\frac{\xi_{max}}{\xi_{min}}} + \sqrt{\frac{\xi_{min}}{\xi_{max}}} \right)^2 \left( \sum_{i=1}^n |\xi_i| \right)^2 \\ nM &\leq \frac{1}{4} \left( \frac{(\xi_{max} + \xi_{min})^2}{\xi_{max} \xi_{min}} \right) [E_{SL^+}(G)]^2 \end{aligned}$$

and inequality (8) follows.  $\square$

**Theorem 3.9.** Let  $G$ ,  $\xi_{max}$ , and  $\xi_{min}$  be same as in Theorem 3.7. Then

$$E_{SL^+}(G) \geq \sqrt{nM - \alpha(n) (\xi_{max} - \xi_{min})^2} \quad (9)$$

where  $\alpha(n)$  is the parameter defined in Theorem 3.5.

*Proof.* Applying Theorem 3.5 for  $a_i = |\xi_i| = b_i$ ,  $a = \xi_{min} = b$ , and  $A = \xi_{max} = B$ , we get

$$\begin{aligned} \left| n \sum_{i=1}^n |\xi_i|^2 - \left( \sum_{i=1}^n |\xi_i| \right)^2 \right| &\leq \alpha(n) (\xi_{max} - \xi_{min})^2 \\ nM - (E_{SL^+}(G))^2 &\leq \alpha(n) (\xi_{max} - \xi_{min})^2 \end{aligned}$$

from which inequality (9) directly follows.  $\square$



Since  $\alpha(n) \leq \frac{n^2}{4}$ , by Theorem 3.9 we get:

**Corollary 3.10.**

$$E_{SL^+}(G) \geq \sqrt{nM - \frac{n^2}{4} (\xi_{max} - \xi_{min})^2}.$$

**Theorem 3.11.** *Let  $G$ ,  $\xi_{max}$ , and  $\xi_{min}$  be same as in Theorem 3.7. Then*

$$E_{SL^+}(G) \geq \frac{M + n \xi_{max} \xi_{min}}{(\xi_{max} + \xi_{min})}. \quad (10)$$

*Proof.* Applying Theorem 3.6 for  $b_i = |\xi_i|$ ,  $a_i = 1$ ,  $r = \xi_{min}$ , and  $R = \xi_{max}$  we get

$$\begin{aligned} \sum_{i=1}^n |\xi_i|^2 + \xi_{max} \xi_{min} \sum_{i=1}^n 1 &\leq (\xi_{max} + \xi_{min}) \sum_{i=1}^n |\xi_i| \\ M + n \xi_{max} \xi_{min} &\leq (\xi_{max} + \xi_{min}) E_{SL^+}(G). \end{aligned}$$

Inequality (10) follows.  $\square$

**Acknowledgment.** The author HSR is thankful to the University Grants Commission (UGC), New Delhi for support through grant under UGC-SAP DRS-III, 2016-2021: F.510/3/DRS-III /2016 (SAP-I).

## References

- [1] N. Abreu, D. M. Cardoso, I. Gutman, E. A. Martins, M. Robbiano, Bounds for the signless Laplacian energy, *Linear Algebra Appl.* **435** (2011) 2365–2374.
- [2] M. Biernacki, H. Pidek, C. Ryll–Nardzewski, Sur une inégalité entre des intégrales définies, *Univ. Maria Curie Skłodowska* **A4** (1950) 1–4.
- [3] B. Borovićanin, K. C. Das, B. Furtula, I. Gutman, Bounds for Zagreb indices, *MATCH Commun. Math. Comput. Chem.* **78** (2017) 17–100.
- [4] D. Cvetković, M. Doob, H. Sachs, *Spectra of Graphs – Theory and Application*, Academic Press, New York, 1980.
- [5] D. Cvetković, P. Rowlinson, S. Simić, Signless Laplacians of finite graphs, *Linear Algebra Appl.* **423** (2007) 155–171.
- [6] D. Cvetković, P. Rowlinson, S. Simić, *An Introduction to the Theory of Graph Spectra*, Cambridge Univ. Press, Cambridge, 2010.

- [7] D. Cvetković, S. K. Simić, Towards a spectral theory of graphs based on the signless Laplacian I, *Publ. Inst. Math. (Beograd)* **85** (2009) 19–33.
- [8] D. Cvetković, S. K. Simić, Towards a spectral theory of graphs based on the signless Laplacian II, *Linear Algebra Appl.* **432** (2010) 2257–2272.
- [9] D. Cvetković, S. K. Simić, Towards a spectral theory of graphs based on the signless Laplacian III, *Appl. Anal. Discr. Math.* **4** (2010) 156–166.
- [10] J. B. Diaz, F. T. Metcalf, Stronger forms of a class of inequalities of G.Pólya–G.Szegő and L.V.Kantorovich, *Bull. Am. Math. Soc.* **69** (1963) 415–418.
- [11] E. Ghorbani, On eigenvalues of Seidel matrices and Haemers conjecture, *Designs, Codes Cryptography* **84** (2017) 189–195.
- [12] R. Grone, R. Merris, The Laplacian spectrum of a graph II, *SIAM J. Discr. Math.* **7** (1994) 221–229.
- [13] R. Grone, R. Merris, V. S. Sunder, The Laplacian spectrum of a graph, *SIAM J. Matrix Anal. Appl.* **11** (1990) 218–238.
- [14] I. Gutman, The energy of a graph, *Ber. Math. Stat. Sect. Forschungsz. Graz* **103** (1978) 1–22.
- [15] I. Gutman, K. C. Das, The first Zagreb index 30 years after, *MATCH Commun. Math. Comput. Chem.* **50** (2004) 83–92.
- [16] I. Gutman, B. Furtula, Survey of graph energies, *Math. Interdisc. Res.* **2** (2017) 85–129.
- [17] I. Gutman, B. Zhou, Laplacian energy of a graph, *Linear Algebra Appl.* **414** (2006) 29–37.
- [18] W. H. Haemers, Seidel switching and graph energy, *MATCH Commun. Math. Comput. Chem.* **68** (2012) 653–659.
- [19] X. Li, Y. Shi, I. Gutman, *Graph Energy*, Springer, New York, 2012.
- [20] R. Merris, Laplacian matrices of graphs: A survey, *Linear Algebra Appl.* **197-198** (1994) 143–176.
- [21] R. Merris, A survey of graph Laplacians, *Linear Multilin. Algebra* **39** (1995) 19–31.
- [22] B. Mohar, The Laplacian spectrum of graphs, in: Y. Alavi, G. Chartrand, O. R. Oellermann, A. J. Schwenk (Eds.), *Graph Theory, Combinatorics and Applications*, Wiley, New York, 1991, pp. 871–898.
- [23] P. Nageswari, P. B. Sarasija, Seidel energy and its bounds, *Int. J. Math. Anal.* **8** (2014) 2869–2871.

- 
- [24] M. R. Oboudi, Energy and Seidel energy of graphs, *MATCH Commun. Math. Comput. Chem.* **75** (2016) 291–303.
- [25] N. Ozeki, On the estimation of inequalities by maximum and minimum values, *J. Coll. Arts Sci. Chiba Univ.* **5** (1968) 199–203 (in Japanese).
- [26] G. Pólya, G. Szegő, *Problems and Theorems in Analysis, Series, Integral Calculus, Theory of Functions*, Springer, Berlin, 1972.
- [27] M. R. Rajesh Kanna, R. P. Kumar, M. R. Farahani, Milovanović bounds for Seidel energy of a graph, *Adv. Theor. Appl. Math.* **10** (2016) 37–44.
- [28] H. S. Ramane, M. M. Gundloor, S. M. Hosamani, Seidel equienergetic graphs, *Bull. Math. Sci. Appl.* **16** (2016) 62–69.
- [29] H. S. Ramane, I. Gutman, M. M. Gundloor, Seidel energy of iterated line graphs of regular graphs, *Kragujevac J. Math.* **39** (2015) 7–12.
- [30] H. S. Ramane, R. B. Jummannaver, I. Gutman, Seidel Laplacian energy of graphs, *Int. J. Appl. Graph Theory* **1** (2017) 74–82.

Harishchandra S. Ramane  
Department of Mathematics,  
Karnatak University,  
Dharwad – 580003, India  
E-mail: hsrामane@yahoo.com

Ivan Gutman  
Faculty of Science,  
University of Kragujevac,  
P.O. Box 60, 34000 Kragujevac, Serbia  
E-mail: gutman@kg.ac.rs

Jayashri B. Patil  
Department of Mathematics,  
Hirasugar Institute of Technology,  
Nidasoshi – 591236, India  
E-mail: js.h@rediffmail.com

Raju B. Jummannaver  
Department of Mathematics,  
Karnatak University,  
Dharwad – 580003, India  
E-mail: rajesh.rbj065@gmail.com